

Reactive to Inert: A Novel Way to Tackle Issues With Chemotherapy

Tushar Ranjan Panda and Malay Patra

Department of Chemical Sciences, Tata Institute of Fundamental Research (TIFR), Mumbai

E-mail: tushar.panda@tifr.res.in , malay.patra@tifr.res.in

Clinically approved platinum anti-cancer drugs cisplatin, carboplatin and oxaliplatin suffer from two major drawbacks: (i) inherent and acquired chemo resistance, that leads to relapse of resistant tumour and (ii) dose limiting toxic side effects such as nephrotoxicity, neurotoxicity and myelosuppression.^{1, 2} Reactivity of platinum (II) drugs correlate linearly to glutathione mediated resistance as well as adverse side effects due to their non-specific biomolecule binding.³ To circumvent these issues, we hypothesized to design substitutionally inert platinum drugs with novel mechanism of action. As positively charged platinum complexes containing planar polypyridyl ligand are capable of binding to *ds*-DNA or proteins through intercalation, a small library of cyclometalated platinum(II) complexes were screened for anticancer properties and a novel lead molecule, compound **4**, was identified which is 25-fold more potent compared to cisplatin in human cervical cancer cell line. Owing to its inertness, the antiproliferative activity of **4** is independent of intracellular glutathione and has been found to overcome platinum resistance in ovarian, lung and prostate cancer cells. While cisplatin and oxaliplatin has plasma half-life of *ca.* 1.5 h, **4** remained unchanged in plasma over 24 h. The *in vivo* efficacy of **4** was confirmed in an A549 xenograft. Importantly, by evaluating the expression levels of inflammatory markers in kidney, we demonstrated that **4** is significantly less nephrotoxic than cisplatin. However, this 1st generation molecule suffers from limited solubility and no cancer targetability. Hence, there stands a need for further improvisation of the molecule to tune its pharmacokinetics. Attaching sugars to platinum drugs have been seen to increase solubility and impart cancer targetability without compromising its toxicity.⁴ Thus, we hypothesized to design sugar conjugates of **4** which we speculate to address the aforementioned issues.

In the presentation, I will be discussing the above mentioned aspects in detail.

References

- (1) Gadre, S.; Manikandan M.; Duari, P.; Chhatar S.; Seetharam, U. K.; and M. Patra, *Chem. Eur. J.*, **2022**, DOI: 10.1002/chem.202201259.
- (2) Johnstone, T. C.; Suntharalingam K; and Lippard, S. J. *Chem. Rev.* **2016**, *116*, 3436–3486.
- (3) Oun, R.; Moussa, Y. E.; and Wheate, N. J. *Dalton Trans.* **2018**, *47*, 6645-6653.
- (4) Patra, M.; Awuah, S.G.; and Lippard, S.J. *J. Am. Chem. Soc.* **2016**, *138*, 12541–12551.